

UT-Komaba-93-04
March 1993

Exponent of n -Ising matter fields coupled to $2d$ gravity

Shinobu Hikami

Department of Pure and Applied Sciences, University
of Tokyo, Meguro-ku, Komaba 3-8-1, Tokyo 153, Japan
and

Laboratoire de Physique Théorique de l'Ecole Normale Supérieure,
24 rue Lhomond, F-75231 Paris Cedex 05, France

Abstract

n - Ising spins on a random surface represented by a matrix model is studied as a model of the 2D gravity coupled to matter field with the central charge $c > 1$. The magnetic field is introduced to discuss the scaling exponent Δ , and the value of this magnetic field exponent is estimated by the series expansion.

The behavior of 2d gravity coupled to a matter field has been studied by a matrix model in the large N limit [1]. The analytic solution by Liouville theory is consistent with the result of the matrix model [2]. However, these exact results are all restricted to the case of the central charge of the matter field $c \leq 1$, and the correct expression for the string susceptibility γ_{st} for the $c > 1$ remains unsolved.

As a matrix model for 2d gravity coupled to matter field of $c > 1$, a multiple n-Ising spin model has been introduced. The series expansion [3,4] and Monte Carlo simulations [5,6,7] have been performed to study the universality of the phase transition. These results suggest the existence of the intermediate region for $c > 1$, where the phase transition remains the second order with nontrivial critical exponents.

In this letter, we extend the perturbational analysis of the previous studies [3,4] by adding the external magnetic field on the Ising spin, and discuss the scaling relation of this multiple n-Ising model coupled to 2d gravity.

The free energy of the one-Ising model coupled to 2d gravity in the presence of the external magnetic field H [8] is given by

$$\begin{aligned} F &= -\ln Z \\ &= -\ln \int dM_+ dM_- \exp[-\text{tr}(\frac{1}{2}M_+^2 + \frac{1}{2}M_-^2 + aM_+M_- - ge^H M_+^4 \\ &\quad - ge^{-H} M_-^4)] \end{aligned} \quad (1)$$

where M_+ and M_- are Hermitian $N \times N$ matrices and they represent the vertices on which the up and down Ising spins are placed respectively. The quantity a is a coupling constant related to the temperature $\beta = 1/kT$ as

$$a = \exp(-2\beta) \quad (2)$$

In the large N limit, the free energy is dominated by the planar diagrams and it has a scaling form near the critical cosmological constant g_c ,

$$F = (g - g_c)^{2-\gamma_{st}} \psi\left(\frac{H}{(g - g_c)^\Delta}\right) \quad (3)$$

We restrict ourselves to the genus zero topology. The coupling constant a has a critical value a_c at which new universal behavior appears. For the one-Ising model, the critical value a_c is $1/4$, and the string susceptibility exponent γ_{st} changes from $-1/2$ of the pure gravity case ($a \neq 1/4$) to $-1/3$ of the Ising case at $a = 1/4$. The exponent γ_{st} and the magnetic field exponent Δ are known for the unitary (p,q) model [9] in which the central charge c is given by $c = 1 - 6(p - q)^2/pq$ for $c < 1$;

$$\gamma_{st} = -\frac{2}{p + q - 1} \quad (4)$$

$$\Delta = (\frac{p}{2} + q)/(p + q - 1) \quad (5)$$

The Ising case corresponds to $p=4$ and $q=3$ ($c=1/2$), and thus we have $\gamma_{st} = -1/3$, $\Delta = 5/6$. There exists a relation between these two exponents, $\Delta = (3 - \gamma_{st})/4$ for $c \leq 1$.

We extend the one-Ising model to the multiple n -Ising model, which is described now by 2^n matrices [3]. Each Ising spin of n -species has up and down two states, and there are 2^n different spin states on the vertices of the diagrams. Each Ising spin has the central charge $1/2$ and total central charge is simply $n/2$, which becomes a central charge of the matter field $c = n/2$.

For the $n=2$ ($c=1$) case, the magnetic field exponent Δ may be obtained from the general expression of the (p,q) model in the limit of $p = q + 1$, $p \rightarrow \infty$ as $\Delta = 3/4$. We will show that indeed this value is obtained by our series expansion.

We use the series expansion about the cosmological constant g for the fixed topology (genus zero), from which the exponents γ_{st} and Δ are extracted by the ratio method. We have already obtained the string susceptibility exponent γ_{st} by the use of the series expansion up to order eight [4]. The coefficients of this series of the m -th derivative of the free energy behaves as

$$\lim_{H \rightarrow 0} d^m F / d^m H = \sum c_k g^k \quad (6)$$

$$c_k \sim A^k k^{-3 + \gamma_{st} + m\Delta} \quad (7)$$

The magnetic field H is applied to one spin component and $(n-1)$ spins remain free of this external magnetic field. The free energy F under this magnetic field is easily calculated in each Feynman diagram. We develop the series expansion up to order eighth including the magnetic field. For example, up to order g^3 , the free energy of n -Ising model with a magnetic field H becomes,

$$\begin{aligned} F = & -2(e^H + e^{-H})g + [16(e^{2H} + e^{-2H} + 2a^2)(1 + a^2)^{n-1} \\ & + 2(e^{2H} + e^{-2H} + 2a^4)(1 + a^4)^{n-1}]g^2 \\ & - \{128[e^{3H} + e^{-3H} + (e^H + e^{-H})(2a^2 + a^4)](1 + 2a^2 + a^4)^{n-1} \\ & + \frac{256}{3}[e^{3H} + e^{-3H} + (e^H + e^{-H})(3a^2)](1 + 3a^2)^{n-1} \\ & + 64[e^{3H} + e^{-3H} + (e^H + e^{-H})(a^2 + 2a^4)](1 + a^2 + 2a^4)^{n-1} \\ & + \frac{32}{3}[e^{3H} + e^{-3H} + (e^H + e^{-H})(3a^4)](1 + 3a^4)^{n-1}\}g^3 \\ & + \dots \end{aligned} \quad (8)$$

where the factor $(1 - a^2)^{-2}$ is absorbed in the redefinition of g .

The quantity A is the inverse of the critical coupling g_c and independent of the number m of the derivative of free energy. Therefore, A should be estimated consistently from the various derivatives of the free energy with respect to the magnetic field, and the scaling relation should be checked with consistent values of the exponents γ_{st} and Δ .

For $n=1$ Ising case, we have an exact value of A as 50.625. This value is different from the exact solution of two matrix model $A = 57.6$ [8,10] due to our modification of the coupling constant g by a factor $(1 - a^2)^2$. We find that our series expansion also gives this critical value and by the method of the extrapolation of the successive ratio of c_k for $k \rightarrow \infty$, we obtain quite accurately the magnetic field exponent $\Delta = 5/6$ from the series of d^2F/d^2H . Once the correct value of A is obtained, the exponents γ_{st} and Δ are accurately estimated in our problem, since the ratio of the coefficients is a monotonic function of the order of degree k . The process of the analysis may be shown by plotting $\Gamma = -(A_c - R_k)k/A_c$ against $1/(k + 1)$ where k is the order of the coupling constant g and the ratio c_k/c_{k-1} is denoted by R_k . This value Γ approaches to the value of $-3 + \gamma_{st} + m\Delta$ for $k \rightarrow \infty$. In fig.1, the analysis of the free energy and the second derivative are represented for $n=1$ Ising model.

For $n=2$ Ising case, there appears a logarithmic singularity. Due to this singularity, the exponent γ_{st} becomes underestimated compared to the correct value $\gamma_{st} = 0$. However, the magnetic field exponent Δ , which is the difference between two exponents of the free energy and the second derivative of the free energy, is not influenced by this logarithmic behavior and it is correctly estimated by the previous ratio method explained before. We have used the values of $a = 1/4$ and $A_c = 54.0$ for $n = 2$ Ising model. By the same method presented in fig.1, we obtain $\Delta = 0.75$ which agrees with the correct value.

For $n=4$ ($c=2$) Ising model, we estimate as $A_c = 59.2$, $a_c = 0.23$ and $\Delta = 0.79$. We have the string susceptibility exponent $\gamma_{st} = 0.04$, which agrees with the previous value [3,4].

For $n=6$ ($c=3$) Ising model, we obtain $\Delta = 0.81$ based on the value $a_c = 0.21$ and $A_c = 63.8$. The exponent γ_{st} is estimated by this method as $\gamma_{st} = 0.1$ which is consistent with the previous result[3,4]. It may be interesting to note that the obtained value of γ_{st} of $c = 3$ appears within the error bar of the value of pure 3d Ising specific exponent α , which has been argued by various methods as 0.11. This coincidence is surprising since our model for $c = 3$ represents the bosonic random surface behavior contrary to the fermionic random surface, which appears in the low temperature expansion of 3d Ising model. If the sign factor due to the fermionic

character has no important effect and can be neglected, this coincidence may be understood [11].

For $n=8$ ($c=4$) Ising model, we take $A_c = 66.0$, $a_c = 0.19$. We have $\Delta = 0.825$ and $\gamma_{st} = 0.2$ for these values.

The critical values of A_c used here are consistent with the previous analysis [4]. The error bar may be ± 0.05 for the estimated value of Δ . The accurate estimation of exponents for $n > 8$ becomes difficult within our eighth order expansion, since the fictitious or ghost second peak of γ_{st} appears for $a > a_c$ as shown in the figures of ref.[4]. There is a tendency that both γ_{st} and Δ are increasing with the increasing value of c . It may be interesting to find the value of c , at which the branched polymer behavior can be seen.

In the following, we discuss the critical behavior near a_c which appears in the large system size, i.e. in the large k limit of the g^k expansion. The coefficient c_k of the free energy in the large k limit becomes the partition function of n -Ising spin on a random surface, and we denote the logarithm of c_k divided by k as $\tilde{f}$,

$$\begin{aligned}\tilde{f} &= -\lim_{k \rightarrow \infty} (\ln c_k)/k \\ &\sim -\ln A(a, H)\end{aligned}\tag{9}$$

This free energy $\tilde{f}$ has a singularity at a_c and following the conventional notations of the critical exponents of the specific heat α and the susceptibility γ , we have

$$\tilde{f} \sim (a - a_c)^{2-\alpha}\tag{10}$$

$$\chi = d^2 \tilde{f} / d^2 H \sim (a - a_c)^{-\gamma}\tag{11}$$

The cosmological constant $A(a, H = 0)$ behaves near the critical value of a_c as

$$A(a, 0) = A(a_c, 0) + C(a - a_c)^{1/\phi}\tag{12}$$

where ϕ is a crossover exponent and it describes the shift of the cosmological constant due to the change of a . The crossover occurs from the pure gravity behavior to n -Ising behavior at a_c . The second derivative of the free energy by a becomes from (9) and (12),

$$d^2 \tilde{f} / d^2 a \sim (a - a_c)^{\frac{1}{\phi}-2},\tag{13}$$

$$\alpha = 2 - \frac{1}{\phi}\tag{14}$$

The exponent ϕ is represented conventionally by $d\nu$, where d is Hausdorff dimension and ν is a critical exponent of the correlation length. Since the specific heat exponent α is written conventionally by $\alpha = 2 - d\nu$, we find

$$1/\phi = d\nu \quad (15)$$

The free energy F has a scaling form for small H and $(a - a_c)$,

$$F = (g - g_c)^{2-\gamma_{st}} \psi\left(\frac{a - a_c}{(g - g_c)^\phi}, \frac{H}{(g - g_c)^\Delta}\right) \quad (16)$$

Therefore, we have a scaling relation between two quantities,

$$|a - a_c|^{1/\phi} \sim H^{1/\Delta} \quad (17)$$

and the magnetization M and the susceptibility χ are expressed by

$$M = d\tilde{f}/dH \sim H^{\frac{1}{\Delta}-1} \sim (a_c - a)^{\frac{1-\Delta}{\phi}} \quad (18)$$

$$\chi = d^2\tilde{f}/d^2H \sim H^{\frac{1}{\Delta}-2} \sim (a - a_c)^{\frac{1-2\Delta}{\phi}} \quad (19)$$

since we have $A(a, 0) = A(a, H) + CH^{1/\Delta}$ and $\tilde{f}$ is given by (9). Thus we find the expressions for the critical exponents α, β and γ in terms of Δ and ϕ which are valid for $c > 1$ case:

$$\alpha = 2 - \frac{1}{\phi}, \beta = \frac{1 - \Delta}{\phi}, \gamma = \frac{2\Delta - 1}{\phi} \quad (20)$$

From these equations, it is possible to compare our estimation of the critical exponents with the Monte Carlo simulations, especially about $\beta/d\nu$ and $\gamma/d\nu$ since these quantities are described by the magnetic field exponent Δ from (15). We note that the previous Monte Carlo results [5,7] for the exponents $\beta/d\nu$, $\gamma/d\nu$ of $c = 1$ does not agree with a correct result.

For $c \leq 1$, we have $\beta = 1/2$ and $\Delta = (3 - \gamma_{st})/4$. Thus the exponent α becomes $\alpha = 2\gamma_{st}/(1 + \gamma_{st})$ [12]. This relation does not hold for $c > 1$ since we have already found that the obtained value of Δ does not satisfy the relation $\Delta = (3 - \gamma_{st})/4$.

The magnetization $M = -d \ln A(a, H)/dH$ is obtained from the calculation of the inverse cosmological constant $A(a, H)$ in our series expansion up to order eight. Indeed we find that the spontaneous magnetization appears below a_c . However it is difficult to extract the accurate value of the exponent β or γ from this spontaneous magnetization. We have obtained rather accurately the value of $\beta/d\nu = 1 - \Delta$ and $\gamma/d\nu = 2\Delta - 1$ through the study of Δ , but the value of $d\nu$ remains unknown.

If the spontaneous magnetization is analyzed more precisely by the higher order expansion, it may be possible to extract the value of $\phi = 1/d\nu$.

We have also tried the specific heat analysis for $n=1$ Ising model based upon the series expansion up to order ten. From the coefficient c_k of the series expansion about g of the free energy F without a magnetic field, the ratio $R_k = c_k/c_{k-1}$ is obtained. Then the quantity A in (7) is estimated by a ratio method as $A_k = (k+1)R_k - kR_{k-1}$. The second derivative of this quantity by a in (2) is proportional to the specific heat. For finite k , there appears a maximum peak around a critical value $a_c = 0.25$ as shown in fig.2. The extrapolation of A_k'' for $k \rightarrow \infty$ is simply done by plotting this value against $1/k$. As shown in fig.2, we have a cusp singularity for the specific heat with an exponent $\alpha = -1$ and a finite size scaling relation $A_\infty'' - A_k'' \sim k^\alpha$. This method seems powerful to extract the exponent $\alpha = 2 - d\nu$, however for $c > 1$, the ghost singularity observed in ref.[4] makes the analysis to be difficult without extending the series expansion higher.

As a summary, we have obtained numerically the value of the scaling exponent of the magnetic field Δ for $c > 1$ by the series analysis of n -Ising model, and we have shown how this exponent is related to other exponents.

The author thank E. Brézin, V. Kazakov and J. Zinn-Justin for the valuable discussions on the present subject. He acknowledges a hospitality of Ecole Normale Supérieure. This work is supported by the Grant-in-Aid for Scientific Research by the Ministry of Education, Science and Culture.

References

- [1] E. Brézin and V. A. Kazakov, Phys. Lett. B236 (1990) 144.
D. J. Gross and A. A. Migdal, Phys. Rev. Lett. 64 (1990) 127.
M. R. Douglas and S. H. Shenker, Nucl. Phys. B335 (1990) 635.
- [2] V. G. Knizhnik, A. M. Polyakov and A. A. Zamolodchikov, Mod. Phys. Lett. A3 (1988) 819.
F. David, Mod. Phys. Lett. A3 (1988) 651.
J. Distler and H. Kawai, Nucl. Phys. B321 (1989) 509..
- [3] E. Brézin and S. Hikami, Phys. Lett. B283 (1992) 203.
- [4] S. Hikami and E. Brézin, Phys. Lett. B295 (1992) 209.
- [5] C. A. Baillie and D. Johnston, Phys. Lett. B 286 (1992) 44.
- [6] S. M. Catterall, J. B. Kogut and R. L. Renken, Phys. Lett. B292 (1992) 277.
- [7] J. Ambjørn, B. Durhuus, T. Jónsson and G. Thorleifsson, NBI-HE-92-35 (1992).
- [8] D. V. Boulatov and V. A. Kazakov, Phys. Lett. B187 (1987) 379.
- [9] B. Eynard and J. Zinn-Justin, Saclay preprint SPhT/93-029.
- [10] M. L. Mehta, Commun. Math. Phys. 49 (1981) 327.
- [11] J. Ambjørn, A. Sedrakyan and G. Thorleifsson, NBI-HE-92-80 (1992).
- [12] V. A. Kazakov, "fields, strings and critical phenomena", Les Houches 1988, edited by E. Brézin and J. Zinn-Justin, P.369, North-Holland, Amsterdam (1990).

figure caption

fig.1 The dots show the values of $\Gamma = -(A_c - R_k)k/A_c$ plotted against $1/(k+1)$, where R_k are ratios of the coefficients in the series of the free energy (upper line) of $n = 1$ Ising model and of the second derivative by a magnetic field (lower line). The intersections of two lines with the vertical axis shows the exponents $3 - \gamma_{st}$ and $3 - \gamma_{st} - 2\Delta$. The value of Δ is obtained as $5/6$.

fig.2 The second derivatives of the inverse cosmological constant $A_k(a)$ by $a = \exp(-2\beta)$ are shown for $k = 6, \dots, 10$ in the $n = 1$ Ising model. The broken line is the extrapolation for $k \rightarrow \infty$ and it shows a cusp singularity at $a = 1/4$.